

Dritsaki, Melina; Dritsaki, Chaido

## Article

# Forecasting European Union CO2 emissions using autoregressive integrated moving average-autoregressive conditional heteroscedasticity models

International Journal of Energy Economics and Policy

## Provided in Cooperation with:

International Journal of Energy Economics and Policy (IJEEP)

*Reference:* Dritsaki, Melina/Dritsaki, Chaido (2020). Forecasting European Union CO2 emissions using autoregressive integrated moving average-autoregressive conditional heteroscedasticity models. In: International Journal of Energy Economics and Policy 10 (4), S. 411 - 423.  
<https://www.econjournals.com/index.php/ijEEP/article/download/9186/5149>.  
doi:10.32479/ijEEP.9186.

This Version is available at:

<http://hdl.handle.net/11159/8436>

## Kontakt/Contact

ZBW – Leibniz-Informationszentrum Wirtschaft/Leibniz Information Centre for Economics  
Düsternbrooker Weg 120  
24105 Kiel (Germany)  
E-Mail: [rights\[at\]zbw.eu](mailto:rights[at]zbw.eu)  
<https://www.zbw.eu/econis-archiv/>

## Standard-Nutzungsbedingungen:

Dieses Dokument darf zu eigenen wissenschaftlichen Zwecken und zum Privatgebrauch gespeichert und kopiert werden. Sie dürfen dieses Dokument nicht für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, aufführen, vertreiben oder anderweitig nutzen. Sofern für das Dokument eine Open-Content-Lizenz verwendet wurde, so gelten abweichend von diesen Nutzungsbedingungen die in der Lizenz gewährten Nutzungsrechte.



<https://zbw.eu/econis-archiv/termsfuse>

## Terms of use:

*This document may be saved and copied for your personal and scholarly purposes. You are not to copy it for public or commercial purposes, to exhibit the document in public, to perform, distribute or otherwise use the document in public. If the document is made available under a Creative Commons Licence you may exercise further usage rights as specified in the licence.*



# Forecasting European Union CO<sub>2</sub> Emissions Using Autoregressive Integrated Moving Average-autoregressive Conditional Heteroscedasticity Models

Melina Dritsaki<sup>1\*</sup>, Chaido Dritsaki<sup>2</sup>

<sup>1</sup>University of Oxford, Oxford, UK, <sup>2</sup>Department of Accounting and Finance, University of Western Macedonia, Kozani, Greece.

\*Email: [melina.dritsaki@ndorms.ox.ac.uk](mailto:melina.dritsaki@ndorms.ox.ac.uk)

Received: 07 January 2020

Accepted: 23 April 2020

DOI: <https://doi.org/10.32479/ijee.9186>

## ABSTRACT

In the past few decades, there are lot of discussions around global warming and climate change primarily due to the increased CO<sub>2</sub> emissions generated by the consumption of fossil fuels, such as oil and natural gas. After an enormous effort, the EU-28 managed to reduce CO<sub>2</sub> emissions in 2014 by 25.7% comparing to 1990 (Kyoto Protocol). This effort should continue in the future so that the EU-28 achieve a 40% reduction on CO<sub>2</sub> emissions by 2030. The current paper aims at investigating the optimum model to forecast CO<sub>2</sub> emissions in the EU-28. To achieve this aim an autoregressive integrated moving average (ARIMA) (1,1,1)- autoregressive conditional heteroscedasticity (ARCH) (1) model was used, combined with the linear ARIMA model and the conditional variance of the ARCH model. The estimation of parameter optimisation of ARIMA(1,1,1)-ARCH(1) model was done with the maximum likelihood approach using the Marquardt (1963), and Berndt-Hall-Hausman algorithms and the three distributions (normal, t-student, generalized error), whereas for the estimation of the covariance coefficient the reversed matrix by Hessian was used. Finally, in order to forecast the ARIMA(1,1,1)-ARCH(1) model, a dynamic as well as a static process was applied. The results of the forecasting revealed that the static procedure provides a better forecast comparing to the dynamic one.

**Keywords:** CO<sub>2</sub> Emissions, Autoregressive Integrated Moving Average (1,1,1)-Autoregressive Conditional Heteroscedasticity (1) Model, Forecasting, E.U

**JEL Classifications:** C22, C53, Q50

## 1. INTRODUCTION

During the last decades, the increase of the greenhouse gas emissions is considered the biggest threat for global warming. The economic growth of developed countries pushes the intensive use of energy and the consumption of fossil fuels, which results to more residues and waste leading to environmental decomposition.

Data from the 1960s and 1970s, show that the concentration of CO<sub>2</sub> in the atmosphere is increasing significantly. Hence, scientists put a lot of pressure to governments to take action.

Unfortunately, it took the international community years to respond to this request. Back in 1988 the International Meteorology Organisation and the United Nations Environmental Program formed an intergovernmental committee for climate change. The evaluation report published in 1990, mentioned that the problem of temperature rise is an existing one and owes to be dealt with promptly. This outcome pushed governments to establish the United Nations Framework Convention on Climate Change (UNFCCC), which was signed in Rio de Janeiro in 1992. UNFCCC as well as the Kyoto Protocol that followed, form the only international frameworks for tackling climate change (United Nations Climate Change, 2019).

The Kyoto Protocol is an international treaty (signed on 11 December 1997, but entered into force on 16 February 2005), which sets the principles of reducing greenhouse gas concentrations in the atmosphere which cause temperature rise in the planet. Based on Kyoto Protocol, countries which have signed the treaty, are bound to reduce greenhouse gas emissions on average by 5.2% between 2008 and 2012, comparing to 1990 levels. The protocol is based on the principle of common responsibilities in tackling climate change but acknowledges the different capacity to achieve this based on each country's economic growth.

Negotiations for resolving the global temperature increase problem were tough due to clashing interests. Consequently, opposing country-teams with diverging views were generated. For example, countries producing carbon, such as Japan, USA, Canada, Australia, New Zealand, but also members of OPEK with Russia and Norway, which support the development of oil and natural gas production, are affected by Kyoto Protocol because they need to reduce their production and instead switch to alternative energy sources. In addition, emerging makers such as China and India could not commit to reducing greenhouse gas. On the contrary, the European Union was the earnest supporter of Kyoto Protocol, which committed to reduce greenhouse gas emissions by 6% for 2012 and reduction of fossil fuel consumption by 16%.

Kyoto Protocol followed a second phase during the period 2013-2020 known as Doha Amendment 2012. Thirty-eight developed countries participate among them 28 member states of the European Union. In this second phase, participating states commit to reduce their emissions to a level 18% lower than that of 1990. The EU also has committed to reduce by 20% for that period.

The Paris agreement for climate change took place in December 2015. This conference set two requirements for its application. The first one concerned the confirmation of the UNFCCC, by at least 55 country-members and the second one concerned the minimum amount of greenhouse gas emissions, which each country need to follow.

The Kyoto Protocol applies to six greenhouse gases: Carbon dioxide (CO<sub>2</sub>), methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), and sulphur hexafluoride (SF<sub>6</sub>). Carbon dioxide (CO<sub>2</sub>) is a natural gas, which is defined by the photosynthesis into organic material. Most of the CO<sub>2</sub> emissions are due to fossil fuel consumption such as carbon, petroleum, natural gas and the burning of biomass. CO<sub>2</sub> emissions are also generated by the change in soil use, the car mobility and various others industrial activities. CO<sub>2</sub> is the main anthropogenic greenhouse gas that affects the radioactive balance of the earth. Moreover, CO<sub>2</sub> is the gas that form the basis in which other greenhouse gas are calculated, resulting in a dynamic overheating of the planet.

The majority of the countries, failed to achieve their commitments with regards to CO<sub>2</sub> emissions. The European Union is taking notable measures for reducing CO<sub>2</sub> emissions generated from the absorption of fossil fuels.

Given that there exists a two-way causal relationship between economic growth and CO<sub>2</sub> emissions, a reduction on the CO<sub>2</sub> emissions will have an unfavourable impact on the economic growth of European Countries Table A1. Table A2, presents the relationship between per capita CO<sub>2</sub> emissions and the per capita economic power of the 28 European Countries.

The amount of CO<sub>2</sub> emissions during the period 1990-2014 in EU, USA, China and the World is presented in the Figure 1.

From the Figure 1 we detect that since the Kyoto Protocol in 1990, the USA have reduced CO<sub>2</sub> emissions by 0.63 % pa and the EU by 1.19% pa. On the contrary, China has increased CO<sub>2</sub> emissions by 5.49% pa whereas the global CO<sub>2</sub> emissions are increasing by 0.73% pa.

The Table 1 presents the descriptive statistics for the per capita CO<sub>2</sub> emissions in the European Union, the USA, China and the World from 1990 (Kyoto Protocol) until 2014. The descriptive statistics mean, standard deviation (Std. Dev.) and coefficient of variation (CV) of these variables are recorded below in Table 1.

Table 1 shows that the variability in the per capita CO<sub>2</sub> emissions is greater in China and smaller in in the case of the USA.

The Figure 2 presents the rate of CO<sub>2</sub> emissions in the EU-28 countries.


The Figure 2 shows an overall downward trend of the CO<sub>2</sub> emissions between 1990 and 1999, with the exception of a peak in 1996, when an exceptional of a cold winter which led to increased demand for heating. From 1999 to 2006, the CO<sub>2</sub> emission for the EU-28 was relative stable. From 2006-2009 a sharp drop of CO<sub>2</sub> emissions was detected as result of the global financial and economic crisis leading to the decrease of industrial activity of the European Union. The CO<sub>2</sub> emissions increased again during the period 2009-2010 and dropped again between 2011 and 2014.

The remainder of the paper is organized as follows: Section 2 provides a brief literature review. Section 3 presents the analysis of methodology. Section 4 summarizes the data. The empirical results are provided in Section 5 and Section 6 proposes the forecasting results. Finally, the last section offers the concluding remarks.


## 2. LITERATURE REVIEW

The forecasting issues are important and are being applied in various scientific fields, such as economics, meteorology, medicine, mechanics, ecology and many more. The increasing trend of the CO<sub>2</sub> emissions in a global level due to human activity indicated the increased atmospheric concentration of CO<sub>2</sub>. Climate change, because of global warming, is one of the most prolific issues during the last years. Reddy et al. (1995) suggest in their research that the total average temperature increase will reach 3-4°C, doubling the CO<sub>2</sub> emissions concentration, whereas in 2007 the intergovernmental committee for climate change reported an increase of the temperature between 1.1 to 6.4°C in the next 100 years.

**Figure 1:** Amount of carbon dioxide emissions (metric tons per capita) from consumption of energy over 1990-2014 in E.U, U.S.A, China and World



**Figure 2:** Amount of carbon dioxide emissions (metric tons per capita) from consumption of energy over 1990-2014 in E.U



**Table 1: Descriptive statistics for CO<sub>2</sub> emissions (1990-2014)**

Variables	Mean	Std. Dev.	CV (%)	Minimum (year)	Maximum (year)
CO <sub>2</sub> E.U	7.880	0.585	7.42	6.379 (2014)	8.540 (1990)
CO <sub>2</sub> USA	18.789	1.209	6.43	16.310 (2012)	20.179 (2000)
CO <sub>2</sub> China	4.173	1.930	46.24	2.152 (1990)	7.557 (2013)
CO <sub>2</sub> World	4.384	0.377	8.59	3.986 (1999)	5.005 (2012)

Developed countries have a higher share of global CO<sub>2</sub> emissions comparing to developing countries. Nakicenovic back in 1994, studies the prospect of greenhouse gas emissions in a rural context. His findings suggest that developing countries are responsible for less than the 16% of the CO<sub>2</sub> concentration due to their previous consumption from mineral sources of energy. Developed countries have a higher share of global emissions. Researchers so far have investigated the forecasting of CO<sub>2</sub> emissions in various countries. Lotfalipour et al. (2013) have examined the economic aspects of CO<sub>2</sub> emissions and their consequences for the case of Iran. In their study they apply Grey and autoregressive integrated moving average (ARIMA) models to forecast CO<sub>2</sub> emission in the period between 1990 and 2011. Their results suggest that Grey models provided better results in terms of forecasting CO<sub>2</sub> emissions. Based on their estimated results, the quantity of CO<sub>2</sub> emissions will reach 925.680.000

tons in 2020, which indicated an increase of 66% compared to 2010, which is highly significant.

Rahman and Hasan (2017), investigated the CO<sub>2</sub> emissions between 1972 and 2015 in the case of Bagladesh. The optimum prognostic model for the CO<sub>2</sub> emissions in the period under investigation was the ARIMA(0, 2, 1) model. The results suggested that the CO<sub>2</sub> emissions for years 2016, 2017 and 2018 will be 83.94657, 89.90464 and 96.28557 metric tons respectively.

Pruethsan in 2007, analysed CO<sub>2</sub> emissions in Thailand using the VARIMAX approach during the period 2000-2015 and subsequently determined the VARIMAX(2, 1, 1) and VARIMAX(2, 1, 3) models as the optimal ones for the CO<sub>2</sub> forecast emissions in Thailand. The forecasting results (using the VARIMAX(2, 1, 1) model) show that CO<sub>2</sub> gas greenhouse

emissions will increase steadily and will reach 25.17% until 2025 comparing to 2016, whereas using the VARIMAX(2, 1, 3) model the CO<sub>2</sub> gas greenhouse emissions will increase steadily and will reach 41.51% until 2040 comparing to 2016.

Nyoni and Mutongi (2019) are using annual data to investigate CO<sub>2</sub> emissions in the case of China from 1960 to 2017 using the Box-Jenkins approach. The ARIMA(1, 2, 1) model was shown to be the most suitable one to forecast CO<sub>2</sub> emissions in the period under investigation. The study results reveal that CO<sub>2</sub> emissions in China will increase and reach approximately 10.000.000 kt in 2024. This result forms a warning of the Chinese government with regards to clinical change and the overheating that China causes in the world.

Finally, Nyoni and Bonga (2019) use 1960-2017 data and a Box-Jenkins approach to forecast CO<sub>2</sub> emissions in China. The study proposes the ARIMA(2, 2, 0) as the optimum one to forecast CO<sub>2</sub> emissions. It further suggests that by 2025, the annual CO<sub>2</sub> emissions in India will reach 3.890.000 kt. The results is critical to the Indian government with respect its short-term and long-term planning of climate change and global overheating.

### 3. THEORETICAL BACKGROUND

#### 3.1. ARIMA Models

An ARIMA model is a generalization of an autoregressive moving average (ARMA) used in econometrics. ARIMA is one of the type of models in the Box and Jenkins (1976) methodology for analysis and forecasting a time series.

The ARIMA( $p, d, q$ ) can be expressed as:

$$\left(1 - \sum_{i=1}^p \phi_i L^i\right) (1-L)^d (y_t - \mu) = \left(1 - \sum_{j=1}^q \theta_j L^j\right) e_t \quad (1)$$

where

$$\phi_p(L) = 1 - \sum_{i=1}^p \phi_i L^i \text{ and } \theta_q(L) = 1 - \sum_{j=1}^q \theta_j L^j \text{ are polynomials}$$

in terms of  $L$  of degree  $p$  and  $q$ .

$y_t$  is the time series, and  $e_t$  is the random error at time period  $t$ , with  $\mu$  is the mean of the model.

$d$  is the order of the difference operator.

$\phi_1, \phi_2, \dots, \phi_p$  and  $\theta_1, \theta_2, \dots, \theta_q$  are the parameters of autoregressive and moving average terms with order  $p$  and  $q$  respectively.

$L$  is the difference operator defined as  $\Delta y_t = y_t - y_{t-1} = (1-L)y_t$ .

#### 3.2. Autoregressive Conditional Heteroscedasticity (ARCH)-GARCH Models

##### 3.2.1. ARCH( $q$ ) model

Engle (1982) developed the ARCH model for testing the volatility of time series. The basic ARCH model consists of two equations,

a conditional mean equation and a conditional variance equation. Both equations form a system that is estimated together with maximum likelihood (ML) method.

So, ARCH model is an ARMA and can be written as follows:

$$y_t = \mu_t + \varepsilon_t \text{ (conditional mean equation)} \quad (2)$$

where  $\mu_t$  is conditional mean of  $y_t$ , and  $\varepsilon_t$  is the shock at time  $t$ .

The variance  $\varepsilon_t$  will be:

$$\varepsilon_t = u_t \sigma_t$$

where  $u_t$  is a white noise with zero mean and variance of one  $u_t \rightarrow iid(0,1)$ .  $u_t$  may or may not follow normal distribution.

$$\sigma_t^2 = \omega + \sum_{i=1}^q \alpha_i \varepsilon_{t-i}^2 \text{ (conditional variance equation)} \quad (3)$$

where  $\sigma_t^2$  is the conditional variance of  $y_t$ ,  $\omega$  is a constant term, and  $q$  is the order of the ARCH terms,  $\omega > 0$ ,  $\alpha_i \geq 0$  and  $i > 0$ .

##### 3.2.2. GARCH( $q, p$ ) model

Bollerslev (1986) extended the ARCH model in a new one that allows the errors of variance to depend on its own lags as well as lags of the squared errors. In other words, it allows the extension of conditional variance to follow an ARMA process.

The GARCH model can be expressed as:

$$y_t = \mu_t + \varepsilon_t \text{ (conditional mean equation)} \quad (4)$$

where  $\mu_t$  is conditional mean of  $y_t$ , and  $\varepsilon_t$  is the shock at time  $t$ .

$$\varepsilon_t = u_t \sigma_t$$

where  $u_t$  is a white noise with zero mean and variance of one  $u_t \rightarrow iid(0,1)$ .

$$\sigma_t^2 = \omega + \sum_{i=1}^q \alpha_i \varepsilon_{t-i}^2 + \sum_{j=1}^p \beta_j \sigma_{t-j}^2 \text{ (conditional variance equation)} \quad (5)$$

where

$\sigma_t^2$  is the conditional variance of  $y_t$ ,  $\omega$  is a constant term,  $q$  is the order of the ARCH terms, and  $p$  is the order of the GARCH terms. We assume that for every  $p \geq 0$  and  $q > 0$ , the parameters are unknown and since the variance is positive, then the following relations must be positive too  $\omega \geq 0$ , and  $\alpha_i \geq 0$  for every  $i=1, \dots, q$  and  $\beta_j \geq 0$  for  $j = 1, \dots, p$ . If the parameters are constrained

such that  $\sum_{i=1}^q \alpha_i + \sum_{j=1}^p \beta_j < 1$ , they imply a weak stationarity. If  $p = 0$ , then GARCH model is becoming an ARCH model.



### 3.3. ARIMA-ARCH/GARCH Model

The ARIMA-ARCH/GARCH model is one model in which the variance of the error term of the ARIMA model follows an ARCH/GARCH process. In other words, the ARIMA-ARCH/GARCH model is a non-linear time series model which combined the lineal model ARIMA with the conditional variance of the ARCH/GARCH model.

For the ARIMA-ARCH/GARCH process to be suggested, the following two phases should be applied. The first one uses the best ARIMA model which fits the stationary and linear data of the time series, whereas the linear model residuals should contain the non-linear part of the data. The second phase uses the ARCH/GARCH model in order to include the non-linear patters of the residuals. The model combined the ARIMA model with the ARCH/GARCH which contains the non-linear patterns of the residuals (Dritsaki, 2018).

The process of parameter estimation of the ARIMA-ARCH/GARCH model is achieved through the logarithmic function of ML through nonlinear least squares using Marquardt's algorithm (1963). The latter is presentd by the following function:

$$\ln L[(y_t), \theta] = \sum_{t=1}^T \left\{ \ln[D(z_t(\theta)), v] - \frac{1}{2} \ln[\sigma_t^2(\theta)] \right\} \quad (6)$$

where  $\theta$  is the vector of the parameters that have to be estimated for the conditional mean, conditional variance and density function,  $z_t$  denoting their density function,  $D(z_t(\theta), v)$ , is the log-likelihood function of  $[y_t(\theta)]$ , for a sample of T observation. The ML estimator  $\hat{\theta}$  for the true parameter vector is found by maximizing (8) (Dritsaki, 2017; 2019).

#### 3.3.1. Conditional Distributions

Logarithmic function of ML used for parameters' estimation on volatility models for all theoretical distributions are the following: (Dritsaki, 2017; 2019).

- Normal distribution

$$\ln L[(y_t), \theta] = -\frac{1}{2} \left[ T \ln(2\pi) + \sum_{t=1}^T z_t^2 + \sum_{t=1}^T \ln(\sigma_t^2) \right] \quad (7)$$

where  $\theta$  is the vector of the parameters that have to be estimated for the conditional mean, conditional variance and density function, T is observations.

- t-student distribution

$$\ln L[(y_t), \theta] = T \left[ \ln \Gamma\left(\frac{v+1}{2}\right) - \ln \Gamma\left(\frac{v}{2}\right) - \frac{1}{2} \ln[\pi(v-2)] \right] - \frac{1}{2} \sum_{t=1}^T \left[ \ln(\sigma_t^2) + (1+v) \ln\left(1 + \frac{z_t^2}{v-2}\right) \right] \quad (8)$$

where

$\Gamma(v) = \int_0^\infty e^{-x} x^{v-1} dx$  is the gamma function and  $v$  is the degree of freedom.

- Generalized error distribution

$$\ln L[(y_t), \theta] = \sum_{t=1}^T \left[ \ln\left(\frac{v}{\lambda}\right) - \frac{1}{2} \left| \frac{z_t}{\lambda} \right|^v - (1+v^{-1}) \ln(2) \right] - \ln \Gamma\left(\frac{1}{v}\right) - \frac{1}{2} \ln(\sigma_t^2) \quad (9)$$

where

$$\lambda = \left[ 2^{-2/v} \frac{\Gamma\left(\frac{1}{v}\right)}{\Gamma\left(\frac{3}{v}\right)} \right]^{1/2}$$

### 3.4. Diagnostic Checking of the Model ARIMA-ARCH/GARCH

Before we accept a fitted model and interpret its findings, it is essential to check whether the model is correctly specified, that is, whether the model assumptions are supported by the data. The diagnostic tests of ARIMA-ARCH/GARCH models are based on residuals. Residuals' normality test is employed with Jarque and Bera (1980) test. Ljung and Box (1978) (Q-statistics) statistic for all time lags of autocorrelation is used for the serial correlation test. Also, for the conditional heteroscedasticity test we use the squared residuals of autocorrelation function.

### 3.5. Forecasting Performance Measures

In order to compare the forecasting performance of ARIMA-ARCH/GARCH models we use the following statistics:

- Mean absolute error (MAE)

$$MAE = \frac{1}{n} \sum_{i=1}^n |Y_i - \hat{Y}_i| \quad (10)$$

$Y_i$  is the vector of observed values of the variable being predicted.

$\hat{Y}_i$  is the vector of n predictions.

It measures the average absolute deviation of forecasted values from original ones.

- Root mean squared error (MSE)

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (Y_i - \hat{Y}_i)^2} \quad (11)$$

- MSE has the following formula

$$MSE = \frac{1}{n} \sum_{i=1}^n (Y_i - \hat{Y}_i)^2 \quad (12)$$

MSE gives an overall idea of the error occurred during forecasting.

- The mean absolute percentage error (MAPE)

$$MAPE = \frac{1}{n} \sum_{i=1}^n \left| \frac{Y_i - \hat{Y}_i}{Y_i} \right| \quad (13)$$

This measure represents the percentage of average absolute error occurred. It is independent of the scale of measurement, but affected by data transformation.

## 4. DATA

Annual data for CO<sub>2</sub> emissions (CO<sub>2</sub>) in the E.U (metric tons per capita), are downloaded from the World Bank's development indicators. The data is for the period from 1960 to 2014.

Figure 3 presents the course of the per capita CO<sub>2</sub> emissions in the European Union between 1960 and 2014 at the level.

Figure 3 shows that the CO<sub>2</sub> emissions at the EU present a random walk. Therefore, we will check for the stationarity of the series and its Figure 4 autocorrelation.

From Figure 4 shows that the auto-regression coefficients decline with rapid pace, which implied that the series is non-stationary.

We then apply the aforementioned tests afresh, in order to investigate the presence of stationarity in the first differences of the series. Figure 5, shows the first differences of the CO<sub>2</sub> emissions.

From Figure 5, we observe that the CO<sub>2</sub> emissions present intense fluctuations in their first differences, which is a possible indication of stationarity. We then test the stationarity with the auto-correlation Figure 6.

Figures 5 and 6 both show that the series is likely to be stationary in its first differences.

The confirmation of the series stationarity is achieved by applying the unit root tests Dickey and Fuller (1979; 1981) and Phillips and Perron (1998).

Figure 3: CO<sub>2</sub> emissions in E.U-28




Figure 4: Auto-regression partial autocorrelation of CO<sub>2</sub> emissions on its level

Sample: 1960 2014 Included observations: 55						
Autocorrelation	Partial Correlation	AC	PAC	Q-Stat	Prob	
		1	0.884	0.884	45.407	0.000
		2	0.770	-0.058	80.441	0.000
		3	0.652	-0.077	106.08	0.000
		4	0.540	-0.047	124.00	0.000
		5	0.440	-0.015	136.15	0.000
		6	0.335	-0.094	143.33	0.000
		7	0.242	-0.026	147.15	0.000
		8	0.141	-0.107	148.48	0.000
		9	0.057	-0.011	148.70	0.000
		10	-0.025	-0.071	148.74	0.000
		11	-0.091	-0.009	149.32	0.000
		12	-0.145	-0.030	150.85	0.000
		13	-0.185	-0.002	153.40	0.000
		14	-0.199	0.049	156.42	0.000
		15	-0.220	-0.075	160.21	0.000
		16	-0.255	-0.130	165.41	0.000
		17	-0.264	0.062	171.18	0.000
		18	-0.283	-0.093	177.97	0.000
		19	-0.275	0.057	184.57	0.000
		20	-0.258	0.005	190.55	0.000
		21	-0.247	-0.053	196.15	0.000
		22	-0.252	-0.119	202.17	0.000
		23	-0.257	-0.022	208.66	0.000
		24	-0.261	-0.051	215.53	0.000

The results of Table 2, confirm that the series is stationary in the first differences. The next step is to determine the ARIMA( $p, q$ ) model, based on the results from Figure 6. The parameters  $p$  and  $q$  of the ARIMA model could be determined from the partial autocorrelation and auto-correlation coefficients comparing them respectively with the critical value  $\pm \frac{2}{\sqrt{n}} = \pm \frac{2}{\sqrt{55}} = \pm 0.262$ .

Moreover, to test for autocorrelation we use the Ljung and Box (1978) test determined by:

$$Q_{LB} = n(n+2) \sum_{k=1}^m \left[ \frac{\hat{\rho}_k^2}{n-k} \right] \sim \chi_m^2 \quad (14)$$

Looking at the values of the partial autocorrelation and autocorrelation coefficients (Figure 6) the value for  $p$  is between  $0 < p < 1$  and respectively, the value for  $q$  will be between  $0 < q < 2$ . Using the values above, we chose the best ARIMA( $p, d, q$ ) model from the lowest values of AIC, SC, and HQ criteria. Table 3 shows the values for  $p$  and  $q$ .

**Table 2: Augmented Dickey–Fuller and Phillips Perron unit root tests**


Variable	ADF		P-P	
	C	C,T	C	C,T
CO <sub>2</sub> EE	-1.700(0)	-1.787(0)	-1.884(4)	-1.786(1)
D CO <sub>2</sub> EE	-5.284(0)*	-6.854(0)*	-5.502(4)*	-6.870(2)*

\*\*\* and \*\* show significant at 1%, 5% and 10% levels respectively. (2) The numbers within parentheses followed by ADF statistics represent the lag length of the dependent variable used to obtain white noise residuals. (3) The lag lengths for ADF equation were selected using Schwarz information criterion. (4) Mackinnon (1996) critical value for rejection of hypothesis of unit root applied. (5) The numbers within brackets followed by PP statistics represent the bandwidth selected based on Newey and West (1994) method using Bartlett Kernel. (6) C=Constant, T=Trend. (7) Δ=First differences

**Table 3: Comparing models using AIC, SIC and HQ tests**

ARIMA model	AIC	SC	HQ
D CO <sub>2</sub> EE			
(1,1,0)	0.167	0.240	0.195
(0,1,1)	0.192	0.265	0.220
(1,1,1)	0.111	0.221	0.153
(0,1,2)	0.170	0.280	0.213
(1,1,2)	0.147	0.294	0.204

**Figure 5: CO<sub>2</sub> emission in their first differences**



**Figure 6: Autocorrelation and partial autocorrelation of CO<sub>2</sub> emissions in their first differences**

Sample: 1960 2014 Included observations: 54								
Autocorrelation		Partial Correlation		AC	PAC	Q-Stat	Prob	
				1	0.263	0.263	3.9539	0.047
				2	0.296	0.244	9.0625	0.011
				3	0.230	0.122	12.188	0.007
				4	0.087	-0.057	12.647	0.013
				5	0.090	-0.004	13.141	0.022
				6	0.234	0.215	16.578	0.011
				7	0.264	0.210	21.080	0.004
				8	0.075	-0.136	21.445	0.006
				9	0.210	0.049	24.421	0.004
				10	0.108	0.036	25.220	0.005
				11	-0.053	-0.137	25.420	0.008
				12	-0.073	-0.189	25.802	0.011
				13	-0.088	-0.118	26.376	0.015
				14	-0.087	0.016	26.950	0.020
				15	0.051	0.155	27.152	0.028
				16	0.067	0.008	27.513	0.036
				17	0.079	0.063	28.018	0.045
				18	-0.105	-0.129	28.944	0.049
				19	0.010	0.101	28.953	0.067
				20	-0.181	-0.106	31.863	0.045
				21	-0.055	0.035	32.142	0.057
				22	-0.070	-0.069	32.602	0.068
				23	-0.003	0.056	32.603	0.088
				24	-0.025	-0.065	32.666	0.111



Results from Table 3, reveal that based on Akaike (AIC), Schwartz (SIC) and Hannan-Quinn (HQ) criteria, ARIMA(1,1,1) model is the most suitable one.

The estimation of the ARIMA(1,1,1) model is achieved by the ML method, whereas the optimisation of the model will be achieved using the Berndt-Hall-Hausman (BHHH), algorithm. The covariance coefficient will be estimated with the inverse Hessian matrix. Table 4, shows the results of estimating the ARIMA(1,1,1) model.

Results from Table 4, show that there aren't any problems with the significance of the coefficients. Also, the coefficient for the error variation estimation, shown as SIGMASQ, is  $\rho_s = 0.058$  and is also statistical significant. The inverse roots of the model are AR = 0.94 and MA = 0.76 and are presented in the following Figure 7.

Figure 7 shows that the inverse roots of the model (inverted AR, MA roots), are within the inverted unit cycle, which confirms that the series under review is stationary.

We then test for heteroscedasticity (ARCH(q)) from the squared residuals of the model above. Table 5 shows the following results.

Table 5 reveals that both auto-correlation coefficients and partial autocorrelation coefficients after the first order, are not statistical significant. Therefore, the ARCH or GARCH procedures should be considered.


## 5. EMPIRICAL RESULTS

Since we found that there exists a first order ARCH-GARCH procedure (Table 5), we could move into the model specification followed by its estimation of the conditional mean and the

**Table 4: Estimation of ARIMA(1,1,1) model**

Dependent Variable: DCO2EE				
Method: ARMA Maximum Likelihood (OPG - BHHH)				
Sample: 1961 2014				
Included observations: 54				
Convergence achieved after 25 iterations				
Coefficient covariance computed using observed Hessian				
Variable	Coefficient	Std. Error	t-Statistic	Prob.
AR(1)	0.936118	0.089612	10.44635	0.0000
MA(1)	-0.758287	0.146692	-5.169241	0.0000
SIGMASQ	0.058056	0.011191	5.187797	0.0000
R-squared	0.160234	Mean dependent var		0.011370
Adjusted R-squared	0.127302	S.D. dependent var		0.265400
S.E. of regression	0.247933	Akaike info criterion		0.111355
Sum squared resid	3.134997	Schwarz criterion		0.221854
Log likelihood	-0.006582	Hannan-Quinn criter.		0.153970
Durbin-Watson stat	2.024627			
Inverted AR Roots	.94			
Inverted MA Roots	.76			

**Figure 7: Inverse roots of autoregressive integrated moving average (1,1,1) model**



conditional variance. Estimation of the ARIMA(1,1,1)-ARCH(1) or ARIMA(1,1,1)-GARCH(1,1) model is achieved with the ML method using the BHHH algorithm, the steps of the Marquardt method (1963), the three distributions (Normal, student's, generalized error), while for the co-variance coefficient the inverse matrix by Hessian is applied. Coefficient estimation as well as residual tests with respect to normality, auto-regression and conditional heteroscedasticity, are presented in Table 6.

From Table 6 shows that the ARIMA(1,1,1)-ARCH(1) model with the GED distribution is the most appropriate one (as it has the highest LogL value). All model coefficients are statistical significant and do not present any issues at the diagnostic issues.

Therefore, we could use this model for forecasting purposes. Table 7 presents the results of the estimation of this model.

The estimate of the regression in the ARIMA(1,1,1)-ARCH(1) model could be presented as:

$$D\ CO_2EE_t = 0.933 \cdot D\ CO_2EE_{t-1} - 0.846e_{t-1} \text{ (conditional mean equation)}$$

$$\sigma_t^2 = 0.018 + 0.956 e_{t-1}^2 \text{ (conditional variance equation)}$$

Figure 8 shows the actual and fitted values of the series, as well as the residuals of the fitted model at the 95% confidence interval.

**Table 5: Autoregressive conditional heteroscedasticity (q) process test**

Sample: 1960 2014 Included observations: 54						
Autocorrelation	Partial Correlation	AC	PAC	Q-Stat	Prob	
		1	0.281	0.281	4.5148	0.034
		2	-0.040	-0.130	4.6093	0.100
		3	-0.209	-0.176	7.1996	0.066
		4	-0.036	0.082	7.2800	0.122
		5	0.175	0.162	9.1653	0.103
		6	0.092	-0.053	9.6966	0.138
		7	0.023	0.026	9.7298	0.204
		8	-0.115	-0.064	10.594	0.226
		9	0.053	0.134	10.782	0.291
		10	-0.071	-0.182	11.124	0.348
		11	-0.026	0.008	11.172	0.429
		12	-0.103	-0.083	11.933	0.451
		13	-0.084	-0.039	12.456	0.491
		14	-0.027	-0.050	12.511	0.565
		15	-0.080	-0.072	13.007	0.602
		16	-0.057	-0.050	13.267	0.653
		17	-0.128	-0.068	14.605	0.624
		18	-0.118	-0.134	15.784	0.608
		19	0.063	0.185	16.130	0.649
		20	0.034	-0.107	16.234	0.702
		21	-0.039	-0.048	16.371	0.748
		22	-0.068	0.018	16.811	0.774
		23	-0.167	-0.176	19.546	0.669
		24	-0.105	-0.081	20.654	0.659

**Table 6: Estimation of ARIMA(1,1,1)-ARCH(1)/GARCH(1,1) models**


Parameter	ARCH(1)			GARCH(1,1)		
	Normal	t-student	GED	Normal	t-student	GED
Mean equation						
AR(1)	0.938*	0.938*	0.932*	0.932*	0.932*	0.924*
MA(1)	-0.839*	-0.839*	-0.845*	-0.864*	-0.864*	-0.873*
Variance equation						
$\omega$	0.019*	0.019*	0.017*	0.012	0.012	0.008
$\alpha$	0.866**	0.866**	0.955*	1.024*	1.022*	1.212*
$\beta$	-	-	-	0.067	0.066	0.069
Persistence	-	-	-	1.091	1.088	1.281
LogL	6.172	DOF=2201	PAR=2.609*	6.330	DOF=635.1	PAR=2.941*
Diagnostic tests						
Q <sup>2</sup> (24)	21.316	21.302	23.714	22.188	22.153	24.806
ARCH-X <sup>2</sup> (1)	0.304	0.303	0.636	0.550	0.542	1.461
Jarque-Bera	1.235	1.236	1.203	1.074	1.074	1.104

\*\*\*, \*\*, \* Show significant at 1%, 5% and 10% levels respectively. (2) ( ) is the order of diagnostic tests. (3) LogL is the value of the logarithmic-likelihood. (4) Q<sup>2</sup>(24) is the Q-statistic of correlogram of squared residuals at twenty-four lags. (5) ARCH-X<sup>2</sup>(1) for autoregressive conditional heteroskedasticity, (6) the persistence is calculated as  $(\alpha_1 + \beta_1)$  for the GARCH model

**Table 7: Estimates of ARIMA(1,1,1)-ARCH(1) model**

Dependent Variable: DCO2EE				
Method: ML ARCH - Generalized error distribution (GED) (OPG - BHHH / Marquardt steps)				
Sample (adjusted): 1962 2014				
Included observations: 53 after adjustments				
Convergence achieved after 30 iterations				
Coefficient covariance computed using observed Hessian				
MA Backcast: 1961				
Presample variance: backcast (parameter = 0.7)				
GARCH = C(3) + C(4)*RESID(-1)^2				
Variable	Coefficient	Std. Error	z-Statistic	Prob.
AR(1)	0.932913	0.033564	27.79469	0.0000
MA(1)	-0.845847	0.048850	-17.31504	0.0000
Variance Equation				
C	0.017907	0.006896	2.596807	0.0094
RESID(-1)^2	0.955557	0.377536	2.531034	0.0114
GED PARAMETER	2.609373	0.831499	3.138156	0.0017
R-squared	0.200663	Mean dependent var		0.008566
Adjusted R-squared	0.184990	S.D. dependent var		0.267131
S.E. of regression	0.241160	Akaike info criterion		-0.057013
Sum squared resid	2.966077	Schwarz criterion		0.128864
Log likelihood	6.510833	Hannan-Quinn criter.		0.014467
Durbin-Watson stat	1.950911			
Inverted AR Roots	.93			
Inverted MA Roots	.85			

**Figure 8:** Actual and fitted values, residuals of the model ARIMA(1,1,1)-ARCH(1) for D CO<sub>2</sub>EE




Looking at Figure 8, the residuals show that there is an ARCH procedure in the data.

## 6. FORECASTING


In order to forecast the ARIMA(1,1,1)-ARCH(1,1) models we use both the dynamic (n-step ahead forecasts) and static (one

step-ahead forecast) procedure. The dynamic procedure computes forecasting for periods after the first sample period, using the former fitted values from the lags of dependent variable and ARMA terms. The static procedure uses actual values of the dependent variable. In the following diagram, we present the criteria for the evaluation of forecasting using the dynamic and static forecast respectively (Dritsaki, 2019).

**Figure 9:** Dynamic and static forecast



**Figure 10:** Time series plot for actual and forecasted D CO<sub>2</sub>EE-28 values



**Table 8:** Forecasted values of E.U-28 CO<sub>2</sub> emissions

Year	D CO <sub>2</sub> EE	Forecasted CO <sub>2</sub> EE	95% Confidence interval	
			Lower	Upper
2015	0.155	6.534	6.490	6.603
2016	-0.235	6.299	6.254	6.341
2017	0.371	6.670	6.627	6.712
2018	-0.564	6.106	6.055	6.172
2019	0.842	6.948	6.912	6.989
2020	-1.270	5.678	5.644	5.703

The forecasted values indicate that the CO<sub>2</sub> present fluctuations until the end of 2020. The great drop as shown by the results of the current study for 2020, is consistent with the commitment which the EU promised by the Kyoto protocol, as well as the amendment of Doha in 2012.

## 7. SUMMARY AND CONCLUSION

The degradation of environment becomes the recrudescence of the environment via the exhaustion of sources such as air, water and soil. This degradation is a consequence of a combination of an already big and constantly growing population, the continuing economic growth, the technological exhaustion of natural resources and the pollutant technology. The results of these damages to the human lifestyle and prosperity have caused a great amount of concern.

Many studies showed that there is an influence of economic growth to environmental degradation. The correlation between



per capita GDP and CO<sub>2</sub> emissions is positive, implying that the increasing per capita GDP leads to increase of CO<sub>2</sub> emissions. No turning point is found at which emissions start to decrease. Market economy mechanisms according to studies' results are not sufficient in the decline of CO<sub>2</sub> emissions. For this reason, legal regulations are required to avoid further environmental degradation.

The purpose of this paper is to model and forecast CO<sub>2</sub> emissions of 28 member countries of EU based on annual data (from 1960 until 2014). Using ARIMA(1,1,1)-ARCH(1) model, ML methodology, Marquardt's algorithm methodology (1963) and BHHH, we forecasted CO<sub>2</sub> emissions for the next 6 years (2015-2020). The results of forecasting showed that CO<sub>2</sub> emissions will display fluctuations until the end of 2020. The year 2020 will present a considerable decrease of CO<sub>2</sub> emissions reaching 33.8% less than the year 1990 (Kyoto Protocol) and will cover by far the commitment of EU countries on the above Treaty. Moreover, after the biblical disasters worldwide (the burning of Amazon forest which covers 60% of the total rainforest, the lack of drinkable water) more countries such as USA, China, India and OPEC countries can adapt with last years' phenomena and decrease CO<sub>2</sub> emissions, avoiding planet's major disaster. According to our examined model, European Union will manage and reduce more CO<sub>2</sub> emissions by 40% in relation to 1990 and once more will be consistent with Kyoto protocol.

## REFERENCES

- Berndt, E., Hall, B., Hall, R., Hausman, J. (1974), Estimation and inference in nonlinear structural models. *Annals of Economic and Social Measurement*, 3(4), 653-665.
- Bollerslev, T. (1986), Generalized autoregressive conditional heteroskedasticity. *Journal of Econometrics*, 31(3), 307-327.
- Box, G.E.P., Jenkins, G.M. (1976), *Time Series Analysis, Forecasting and Control*. San Francisco: Holden-Day.
- Dickey, D.A., Fuller, W.A. (1979), Distributions of the estimators for autoregressive time series with a unit root. *Journal of American Statistical Association*, 74(366), 427-431.
- Dickey, D.A., Fuller, W.A. (1981), Likelihood ratio statistics for autoregressive time series with a unit root. *Econometrica*, 49(4), 1057-1072.
- Dritsaki, C. (2017), An empirical evaluation in GARCH volatility modeling: Evidence from the Stockholm stock exchange. *Journal of Mathematical Finance*, 7, 366-390.
- Dritsaki, C. (2018), The performance of hybrid ARIMA-GARCH modelling and forecasting oil price. *International Journal of Energy Economics and Policy*, 8(3), 14-21.
- Dritsaki, C. (2019), Modeling the volatility of exchange rate currency using GARCH model. *Economia Internazionale*, 2(2), 209-230.
- Engle, F.R. (1982), Autoregressive conditional heteroscedasticity with estimates of variance of United Kingdom inflation. *Econometrica*, 50, 987-1008.
- Jarque, C., Bera, A. (1980), Efficient tests for normality, homoscedasticity and serial independence of regression residuals. *Economics Letters*, 6, 255-259.
- Ljung, G.M., Box, G.E.P. (1978), On a measure of a lack of fit in time series models. *Biometrika*, 65(2), 297-303.
- Lotfalipour, M.R., Falahi, M.A., Bastam, M. (2013), Prediction of CO<sub>2</sub> emissions in Iran using grey and ARIMA models. *International Journal of Energy Economics and Policy*, 3(3), 229-237.
- MacKinnon, J.G. (1996), Numerical distribution functions for unit root and cointegration tests. *Journal of Applied Econometrics*, 11(6), 601-618.
- Marquardt, D.W. (1963), An algorithm for least squares estimation of nonlinear parameters. *Journal of the Society for Industrial and Applied Mathematics*, 11, 431-441.
- Nakicenovic, N. (1994), *Energy Strategies for Mitigating Global Change. Technology Responses to Global Environmental Challenges*. Paris: OECD/IEA. p449-459.
- Newey, W.K., West, K.D. (1994), Automatic lag selection in covariance matrix estimation. *Review of Economic Studies*, 61(4), 631-653.
- Nyoni, T., Bonga, W.G. (2019), Prediction of CO<sub>2</sub> emissions in India using ARIMA models, *Dynamic Research Journals*, 4(2), 1-10.
- Nyoni, T., Mutongi, C. (2019), Modeling and Forecasting Carbon Dioxide Emissions in China using Autoregressive Integrated Moving Average (ARIMA) Models. MPRA Paper, No. 93984.
- Phillips, P.C.B., Perron, P. (1998), Testing for a unit root in time series regression. *Biometrika*, 75(2), 335-346.
- Pruethsan, S. (2017), VARIMAX model to forecast the emission of carbon dioxide from energy consumption in rubber and petroleum industries sectors in Thailand. *Journal of Ecological Engineering*, 18(3), 112-117.
- Rahman, A., Hasan, M.M. (2017), Modeling and forecasting of carbon dioxide emissions in Bangladesh using Autoregressive Integrated Moving Average (ARIMA) models. *Open Journal of Statistics*, 7, 560-566.
- Reddy, V.R., Reddy, K.R., Acock, B. (1995), Carbondioxide and temperature interactions on stem extension, node initiation and fruiting in cotton. *Agriculture, Ecosystems and Environment*, 55, 17-28.
- United Nations Climate Change. (2019), Conference 25<sup>th</sup> Website. New York, United States: United Nations Climate Change.



## APPENDIX

**Table A1: CO<sub>2</sub> emissions (metric tons per capita)  
European Union, United States, China, World**

ETH	E.U	U.S.A	China	World
1960	5.765	16.000	1.170	3.099
1961	5.925	15.681	0.836	3.070
1962	6.213	16.014	0.661	3.141
1963	6.578	16.482	0.640	3.245
1964	6.794	16.968	0.626	3.361
1965	6.966	17.452	0.666	3.440
1966	7.119	18.121	0.711	3.539
1967	7.227	18.598	0.574	3.578
1968	7.594	19.089	0.605	3.684
1969	8.008	19.858	0.725	3.824
1970	8.492	21.111	0.943	4.015
1971	8.658	20.980	1.042	4.074
1972	8.946	21.749	1.081	4.158
1973	9.348	22.511	1.098	4.299
1974	9.162	21.503	1.097	4.224
1975	8.944	20.402	1.250	4.121
1976	9.485	21.158	1.285	4.285
1977	9.411	21.532	1.389	4.343
1978	9.680	21.973	1.529	4.320
1979	10.053	21.780	1.543	4.482
1980	9.733	20.786	1.495	4.358
1981	9.156	19.767	1.460	4.150
1982	8.929	18.590	1.567	4.041
1983	8.780	18.572	1.629	3.954
1984	8.702	18.977	1.750	4.025
1985	8.904	18.882	1.871	4.074
1986	8.859	18.721	1.939	4.124
1987	9.002	19.350	2.038	4.152
1988	8.856	20.010	2.151	4.227
1989	9.080	20.076	2.153	4.244
1990	8.584	19.323	2.152	4.194
1991	8.583	19.056	2.229	4.173
1992	8.312	19.139	2.309	4.068
1993	8.171	19.347	2.443	4.002
1994	8.039	19.361	2.566	4.011
1995	8.153	19.277	2.756	4.036
1996	8.362	19.496	2.844	4.071
1997	8.157	19.690	2.821	4.082
1998	8.156	19.579	2.677	4.050
1999	7.992	19.727	2.649	3.986
2000	8.013	20.179	2.697	4.038
2001	8.175	19.637	2.742	4.081
2002	8.106	19.613	3.007	4.088
2003	8.225	19.564	3.524	4.258
2004	8.210	19.658	4.038	4.414
2005	8.145	19.592	4.523	4.528
2006	8.179	19.094	4.980	4.636
2007	8.010	19.218	5.335	4.671
2008	7.804	18.462	5.702	4.762
2009	7.157	17.158	6.010	4.662
2010	7.356	17.443	6.561	4.835
2011	7.079	16.977	7.242	4.975
2012	6.918	16.310	7.425	5.005
2013	6.754	16.323	7.557	4.998
2014	6.379	16.503	7.544	4.981

Source: World development indicators

**Table A2: States in Europe by GDP (PPP) per capita, and  
CO<sub>2</sub> emissions**

Country	CO <sub>2</sub> emissions (per capita)	GDP (PPP) per capita \$
European Union	6.379	52.550
Austria	6.870	46.778
Belgium	8.328	43.338
Bulgaria	5.872	18.292
Croatia	3.974	21.240
Cyprus	5.260	32.373
Czech Republic	9.166	30.433
Denmark	5.936	46.223
Estonia	14.849	27.856
Finland	8.661	40.771
France	4.573	40.801
Germany	8.889	46.627
Greece	6.180	26.017
Hungary	4.266	25.553
Ireland	7.314	52.133
Italy	5.271	35.310
Latvia	3.498	23.487
Lithuania	4.378	27.537
Luxembourg	17.400*	99.738*
Malta	5.401	34.921
Netherlands	9.920	48.363
Poland	7.517	25.334
Portugal	4.332	27.218
Romania	3.498**	19.855**
Slovak Republic	5.662	28.641
Slovenia	6.214	29.879
Spain	5.034	33.285
Sweden	4.478	46.410
United Kingdom	6.497	40.762
World	4.981	15.332

Source: World development indicators, \*Countries with large amount of CO<sub>2</sub> emissions, and high GDP (PPP) (per capita), \*\*Countries with small amount of CO<sub>2</sub> emissions, and low GDP (PPP) (per capita)